

NONABELIAN NATURE OF ASYMMETRY FOR THE B_c MESON PRODUCTION IN GLUON-PHOTON INTERACTION

A.V.Berezhnoy, V.V.Kiselev, A.K.Likhoded

Institute for High Energy Physics,
Protvino, Moscow Region, 142284, Russia

E-mail: kiselev@mx.ihep.su Fax: +7-(095)-2302337

Abstract

Calculations of cross-sections for the gluon-photon production of B_c and B_c^* mesons as well as for the production in ep -interactions at HERA are performed in the leading order of perturbative QCD. We show that the nonabelian structure of QCD leads to an essential forward-backward asymmetry in these processes.

1 Introduction

A significant progress in the understanding of the mechanisms for the production of heavy quarkonium consisting of two heavy quarks with different flavors, for instance, $(\bar{b}c)$ -system, has been achieved during past three years [1, 2, 3, 4]. One clarified that for the B_c production in e^+e^- -annihilation, the differential cross-section has the simple factorized form in the limit of high energies ($M_{B_c}^2/s \ll 1$) [5],

$$\frac{d^2\sigma_{B_c}}{dzdp_T} = \left. \frac{d\sigma_{b\bar{b}}}{dk_T} \right|_{k_T=\frac{p_T}{z}} \cdot \frac{D_{\bar{b} \rightarrow B_c}(z)}{z}, \quad (1)$$

where $z = 2E_{B_c}/\sqrt{s}$, $\sigma_{b\bar{b}}$ is the Born cross-section for the $b\bar{b}$ -pair production, k_T is the transverse momentum of $\bar{b}$ -quark, p_T is the transverse momentum of B_c meson, and $D_{\bar{b} \rightarrow B_c}(z)$ is interpreted as the fragmentation function of $\bar{b} \rightarrow B_c + X$. In the language of Feynmann diagrams, the latter fact means that in the high energy limit one can conveniently choose a gauge, namely the axial gauge, with the unit n vector directed along the four-momentum of the b -quark, so that the contribution of the diagrams where the $c\bar{c}$ -pair is produced off the $\bar{b}$ -quark line is dominant. Using this fact, one analytically calculated the fragmentation functions for the bound states in S -, P - and D -waves [5].

The production of $(\bar{b}c)$ -quarkonium in hadronic, photonic and photon-gluon interactions occurs in a more complex way and it can not be described by eq. (1) in the entire kinematical region even at high energies $M_{B_c}^2/s \ll 1$. The number of diagrams for the mentioned processes is greater than that in e^+e^- -annihilation (one has 36 diagrams for the gluonic interactions, 20 for the photonic collisions and 24 for the photon-gluon subprocess). So, one can find no gauge to restrict the number of dominant diagrams by those of the fragmentation kinds, i.e., by diagrams like 15 or 18 in Fig. 1, where the tree process $g\gamma \rightarrow B_c + X$ is presented. The origin of the problem is that the recombination diagrams also contribute to the cross-section, for example, diagram 5 in Fig. 1 is of this kind. This contribution is essential up to the transverse momentum values equal to 35 GeV. At $p_T > 35$ GeV, the regime of fragmentation

(1) becomes dominant, as one must expect by the theorem on the hard process factorization [6]. However, the latter fact underlines the calculations of the contributions, determined by the complete set of diagrams. The complete perturbative calculations can lead to a possible interference of various contributions in some kinematical regions. So, detailed analysis for the gluonic production of B_c mesons at high energies was performed by us [7]. We found that the effect of destructive interference between the fragmentation diagrams and the ones containing the three-gluon vertex is important, so that in the region of z close to 1 and at moderate p_T values the complete set of diagrams leads to smaller cross-section in comparison with the one calculated according to (1) [7]. We also found that for the photon-photon interaction the interference is absent. These investigations lead us to the conclusion that destructive interference takes place if the interaction possesses the nonabelian character. Therefore, it is interesting to study the production of B_c meson in the gluon-photon interactions, wherein the mentioned interference would result in the forward-backward asymmetry, which is described in this paper.

Unfortunately, a straightforward comparison of experimental measurements with the fragmentation picture is problematic since the condition $M_{B_c}^2/s \ll 1$ is generally not satisfied in partonic subprocesses, and the dominant contribution to the cross-section is integrated over the region close to the threshold of the meson production in the subprocess. In this work the cross-section of B_c production in the HERA experiments will be presented. Despite the dominance of the kinematical region near the subprocess threshold, the effect caused by the presence of the three-gluon vertices, i.e. the nonabelian texture of QCD, turns out to be significant.

2 Calculation technique

The amplitude for the B_c meson production, A^{SJj_z} , can be expressed through the amplitude of four free quarks production, $T^{Ss_z}(p_i, k(\mathbf{q}))$, and the orbital wave function of the B_c meson, $\Psi^{Ll_z}(\mathbf{q})$, in the meson rest frame as

$$A^{SJj_z} = \int T^{Ss_z}(p_i, k(\mathbf{q})) \cdot (\Psi^{Ll_z}(\mathbf{q}))^* \cdot C_{s_z l_z}^{J j_z} \frac{d^3 \mathbf{q}}{(2\pi)^3}, \quad (2)$$

where J and j_z are the total spin of the meson and its projection on z axis in the B_c rest frame, respectively; L and l_z are the orbital momentum and its projection; S and s_z are the sum of quark spins and its projection; $C_{s_z l_z}^{J j_z}$ are the Clebsch-Gordan coefficients; p_i are four-momenta of B_c , b and $\bar{c}$, $\mathbf{q}$ is the three-momentum of $\bar{b}$ quark in the B_c meson rest frame; $k(\mathbf{q})$ is the four-momentum, obtained from the four-momentum $(0, \mathbf{q})$ by the Lorentz transformation from the B_c rest frame to the system, where the calculation of $T^{Ss_z}(p_i, k(\mathbf{q}))$ is performed. Then, the four-momenta of $\bar{b}$ and c quarks, forming the B_c meson, will be determined by the following formulae with the accuracy up to $|\mathbf{q}|^2$ terms

$$\begin{aligned} p_{\bar{b}} &= \frac{m_b}{M} P_{B_c} + k(\mathbf{q}), \\ p_c &= \frac{m_c}{M} P_{B_c} - k(\mathbf{q}), \end{aligned} \quad (3)$$

where m_b and m_c are the quark masses, $M = m_b + m_c$, and P_{B_c} is the B_c momentum. Let us note that for the S -wave states it is enough to take into account only terms independent of $\mathbf{q}$.

The 24 diagrams, contributing to the gluon-photon production of B_c are shown in Fig. 1. The product of spinors $v_{\bar{b}}\bar{u}_c$, corresponding to the $\bar{b}$ and c quarks in the amplitude $T^{Ss_z}(p_i, k(\mathbf{q}))$ of eq.(2), should be substituted by the projection operator

$$\mathcal{P}(\Gamma) = \sqrt{M} \left(\frac{\frac{m_b}{M} \hat{P}_{B_c} + \hat{k} - m_b}{2m_b} \right) \Gamma \left(\frac{\frac{m_c}{M} \hat{P}_{B_c} - \hat{k} + m_c}{2m_c} \right), \quad (4)$$

where $\Gamma = \gamma^5$ for $S = 0$, or $\Gamma = \varepsilon^*(P_{B_c}, s_z)$ for $S = 1$, where $\varepsilon(P_{B_c}, s_z)$ is the polarization vector for the spin-triplet state.

The fact that the $\bar{b}c$ -system is produced in the colour-singlet state simplifies the texture of colour matrix. The latter has the only nonzero eigenvalue for the given process, so that one can significantly reduce the time for the computation of the matrix element.

The S -wave production amplitude can be written down as

$$A^{Ss_z} = iR_S(0) \sqrt{\frac{2M}{2m_b 2m_c}} \sqrt{\frac{1}{4\pi}} \left(T^{Ss_z}(p_i, k(\mathbf{q} = 0)) \right), \quad (5)$$

where $R_S(0)$ is the radial wave function at the origin, so that $\Psi(0) = R(0)/\sqrt{4\pi}$ and

$$R_S(0) = \sqrt{\frac{\pi M}{3}} \tilde{f}_{B_c},$$

and the quantity $\tilde{f}_{B_c}$ is related to the leptonic decay constants of pseudoscalar and vector B_c states

$$\begin{aligned} \langle 0 | J_\mu(0) | V \rangle &= i f_V M_V \epsilon_\mu, \\ \langle 0 | J_{5\mu}(0) | P \rangle &= i f_P p_\mu, \end{aligned}$$

where $J_\mu(x)$ and $J_{5\mu}(x)$ are the vector and axial-vector currents of the constituent quarks. In the lowest order in α_s one has $\tilde{f} = f_V, f_P$. The account for hard gluon corrections in the first order in α_s [8, 9] results in

$$\tilde{f} = f_V \left[1 - \frac{\alpha_s^H}{\pi} \left(\frac{m_2 - m_1}{m_2 + m_1} \ln \frac{m_2}{m_1} - \frac{8}{3} \right) \right], \quad (6)$$

$$\tilde{f} = f_P \left[1 - \frac{\alpha_s^H}{\pi} \left(\frac{m_2 - m_1}{m_2 + m_1} \ln \frac{m_2}{m_1} - 2 \right) \right], \quad (7)$$

where $m_{1,2}$ are the masses of the quarks composing the quarkonium. For the vector currents of quarks with equal masses, the BLM procedure of scale fixing in the "running" coupling constant of QCD [10] gives (see ref. [11])

$$\alpha_s^H = \alpha_s^{\overline{\text{MS}}}(e^{-11/12} m_Q^2).$$

The estimates of the $\tilde{f}_{B_c}$ value within the potential models and QCD sum rules are somewhat uncertain, $\tilde{f}_{B_c} = 460 \pm 100$ MeV [2]. Recent estimates within the QCD sum rules and in the framework of lattice computations give $f_{B_c} = 385 \pm 25$ MeV and $f_{B_c} = 395(2)$ MeV, respectively. Taking into account radiative corrections, the value of $\tilde{f}_{B_c}$ is about 470 MeV [12], which will be used in the following estimates. Since the cross-section is proportional to $\tilde{f}_{B_c}^2$, the corresponding value of the cross-section for a different value of $\tilde{f}_{B_c}$ can be obtained by rescaling with the multiplicative factor $(\tilde{f}_{B_c}/\tilde{f}'_{B_c})^2$.

3 Results of calculations

In the leading order of perturbation theory, the total cross-section for the photon-gluon production of B_c is proportional to $\sigma \sim \frac{\alpha\alpha_s^3}{m_b^2 m_c^2} |\Psi(0)|^2$, where $|\Psi(0)|$, α_s and masses of quarks are free parameters of the model. The calculations are performed with the following values of the parameters

$$\begin{aligned}\alpha &= 1/133, \\ \alpha_s &= 0.24, \\ m_b &= 4.8 \text{ GeV}, \\ m_c &= 1.5 \text{ GeV}, \\ \tilde{f}_{B_c} &= 470 \text{ MeV}.\end{aligned}\tag{8}$$

The results for the gluon-photon production of S -wave levels in the $(\bar{b}c)$ -quarkonium (B_c and B_c^*) are presented in Fig. 2 and Tab. 1. For the sake of comparison, the predictions of the fragmentation mechanism are also shown in Fig. 2. One can see that the fragmentation mechanism underestimates the cross-section, much the same way as in the photonic and gluonic interactions producing B_c (or B_c^*) [3, 4].

The distributions $d\sigma/dz$ for both the B_c meson moving in the gluon hemisphere (i.e. at $p_z > 0$, where p_z is the projection of the momentum on the beam axis in the c.m.s. of the colliding particles) and B_c moving in the photon hemisphere, are shown in Fig. 3. According to the fragmentation model, the mesons are produced symmetrically in both the hemispheres, so that the distributions over z are described by the expression $D(z) \cdot \sigma_{b\bar{b}}/2$ at $p_z > 0$ as well as at $p_z < 0$. However, one can see that the real picture of production is more complex. First, the large enhancement compared to the predictions of the fragmentation model is found in the region of low z for both the hemispheres. Second, a significant asymmetry takes place for $z > 0.5$. Both effects are caused by the following reasons. As was shown in [13], the first reason is the contribution of diagrams like the diagram 20 in Fig. 1. In the leading order of perturbation theory the latter contribution corresponds to the "resolved" photon split into the pair of charmed quarks, so that one of those recombines with the produced $\bar{b}$ -quark to form the B_c -meson. Taking into the ratio of the quark charges $(Q_c/Q_b)^2 = 4$, this mechanism causes the increase of B_c -production cross-section in the photon hemisphere. The second reason is the destructive interference between the $\bar{b}$ -quark fragmentation and the contribution caused by the three-gluon vertices appearing in the splitting of initial gluon inside the cone of the latter (see diagrams 13, 14, 17, 18, 22 and 3, 4). This is why the essential decrease of B_c -production cross-section occurs in the gluon hemisphere.

The distributions $d\sigma/dp_T$ for the forward and backward hemispheres ($p_z > 0$ and $p_z < 0$, respectively) are presented in Fig. 4 for the B_c state. One can see that these distributions differ from the ones obtained in the fragmentation model at low transverse momenta. In the region of high p_T , where the fragmentation mechanism dominates, the deviation becomes small, as one expects.

Comparing Figs. 3 and 4, one can draw the conclusion that the difference between the production cross-sections in the two hemispheres is most significant at large z and low p_T . Our calculations show that the same conclusion is valid for the production of the excited vector state B_c^* .

The considered interactions between the monochromatic beams of gluons and photons do not occur in a real experiment. To clarify whether one can observe the asymmetry in ep or γp

experiments or not, the process of ep -interaction at HERA ($\sqrt{s} = 314$ GeV) is considered. For this purpose, the cross-section of the gluon-photon subprocess has been convoluted with the structure functions of both the photon in electron and the gluon in proton, respectively. The equivalent photon parameterization in [14] has been chosen as the photonic structure function. The CTEQ4 functions have been used for the gluon distributions [15].

The rapidity distribution normalized to unity is shown in Fig. 5 for the production of B_c^* . For comparison, we have also presented the distribution with a constant matrix element, $|A|^2 = \text{const.}$ One can see that the differential cross-section for B_c^* production is strongly displaced in comparison with the distribution calculated with the constant matrix element. Therefore, the effect of destructive interference, which has been found at high energies of the $g\gamma \rightarrow B_c(B_c^*) + X$ subprocess, survives in the real conditions of interaction at HERA, too. On the other hand, in the fragmentation regime at high p_T the asymmetry must tend to zero. The latter practically occurs in Fig. 6, wherein the cross-section $d\sigma/dy$ at $p_T > 30$ GeV is shown in comparison with the distribution for the constant matrix element at the same cut off, $p_T > 30$ GeV.

Summarizing the results above, one can conclude that the effect of the forward-backward asymmetry in the $B_c^{(*)}$ -meson distribution $d\sigma/dy$ is basically related with the nonabelian texture of QCD, and this asymmetry is determined by the three-gluon vertex.

As has been mentioned above, the total cross-section of B_c and B_c^* production at the HERA facilities is about to 2.5 pb including charge-conjugate states. The vector B_c^* state decays to the ground B_c meson through the emission of a γ -quantum in the M1-radiative transition, which has the branching ratio close to 100% [2]. So, at the HERA luminosity $\mathcal{L} = 16 \cdot 10^{30} s^{-1} cm^{-2}$ the calculated value of the cross-section will give only 400 events with the B_c meson per year. The increase of total energy by three times, which correspond to the combination of LEP and LHC accelerators, will result in the enhancement of the asymmetry as well as the total cross-section, which is estimated to have a value of 6 nb at $\sqrt{s} = 660$ GeV (see Fig. 7). At the luminosity $10^{32} s^{-1} cm^{-2}$ one can expect about $6 \cdot 10^3$ $B_c^{(*)}$ events per year.

Taking into account low branching fractions for the exclusive decay modes of B_c (for example, $Br(B_c^+ \rightarrow J/\psi e^+ \nu) \simeq 2.7\%$) and the efficiency for the detecting of decay process ($\epsilon \simeq 0.1$), one can see that the expected number of events can turn out to be insufficient for the investigation of the described asymmetry in the ep -production of B_c meson.

The study of these nonabelian effects is a more real problem for the production of D_s meson. As we can see above, the asymmetry is caused by quite similar deviation from the fragmentation mechanism due to the process-dependent contributions, which have a more rapid decrease with the growth of the transverse momentum of the produced meson. For the $B_c^{(*)}$ mesons these sub-leading terms can be reliably calculated in perturbative QCD. In the D_s production, higher orders in α_s can be significant. Nevertheless, we believe that the sub-leading non-fragmentational contributions have the same feature as in the B_c production: the yield in the photon hemisphere is enhanced and the one in the gluon hemisphere is suppressed due to the nonabelian interactions of gluons. One has to note that the quark-mass ratio determining the asymmetry value in the model under consideration is one and the same for the D_s and B_c mesons, and hence the effect of destructive interference can be experimentally observed at HERA, where the energy scale is given by the ratio of $M_{D_s}/\sqrt{s}$, which is approximately equal to $M_{B_c}/\sqrt{s}$ at the energy of LEP and LHC combination. The distribution for the D_s -meson rapidity at the HERA energy is shown in Fig. 8.

A more detailed study of the interference effect requires the investigation of the contribu-

tions by individual diagrams in a proper gauge. The latter consideration will be presented elsewhere.

This work is in part supported by the Russian Foundation for Basic Research, grants 96-02-18216 and 96-15-96575. V.V.K. and A.K.L. express their gratitude to Profs.A.Wagner and P.M.Zerwas for kind hospitality and support during the visit to DESY, where this work was completed.

References

- [1] S.S.Gershtein, V.V.Kiselev, A.K.Likhoded, A.V.Tkabladze, *Uspekhi Fiz. Nauk* **165**, 3 (1995).
- [2] E.Eichten, C.Quigg, *Phys. Rev.* **D49**, 5845 (1994);
S.S.Gershtein, V.V.Kiselev, A.K.Likhoded, A.V.Tkabladze, *Phys. Rev.* **D51**, 3613 (1995).
- [3] K.Kolodziej, A.Leike, R.Rückl, *Phys. Lett.* **B355**, 337 (1995);
A.V.Berezhnoy, V.V.Kiselev, A.K.Likhoded, *Z. Phys.* **A356**, 79 (1996).
- [4] A.V.Berezhnoy, A.K.Likhoded, M.V.Shevlyagin, *Phys. Lett.* **B342**, 351 (1995);
K.Kolodziej, A.Leike, R.Rückl, *Phys. Lett.* **B348**, 219 (1995);
A.V.Berezhnoy, V.V.Kiselev, A.K.Likhoded, *Phys. Lett.* **A356**, 341 (1996);
A.V.Berezhnoy, V.V.Kiselev, A.K.Likhoded, *Z. Phys.* **A356**, 89 (1996).
- [5] E.Braaten, K.Cheung, T.C.Yuan, *Phys. Rev.* **D48**, 4230 (1993);
V.V.Kiselev, A.K.Likhoded, S.M.Shevlyagin, *Z. Phys.* **C63**, 77 (1994);
T.C.Yuan, *Phys. Rev.* **D50**, 5664 (1994);
K.Cheung, T.C.Yuan, *Phys. Rev.* **D53**, 3591 (1996).
- [6] J.C.Collins, D.E.Soper, *Ann. Rev. Nucl. Part. Sci.* **37**, 383 (1987).
- [7] A.V.Berezhnoy, V.V.Kiselev, A.K.Likhoded, *Yad. Fiz.* **60**, 353 (1997).
- [8] E.Braaten, S.Fleming, *Phys. Rev.* **D52**, 181 (1995).
- [9] V.V.Kiselev, *Int. J. Mod. Phys.* **A11**, 3689 (1996).
- [10] S.J.Brodsky, G.P.Lepage, P.B.Mackenzie, *Phys. Rev.* **D28**, 228 (1983).
- [11] M.Voloshin, *Int. J. Mod. Phys.* **A10**, 2865 (1995).
- [12] V.V.Kiselev, *Phys. Lett.* **B372**, 326 (1996).
- [13] A.V.Berezhnoy, V.V.Kiselev, A.K.Likhoded, *Yad. Fiz.* **59**, 2032 (1996).
- [14] R.M.Barnett et al., PDG, *Phys. Rev.* **D54**, 1 (1996).
- [15] H.L.Lai et al., CTEQ Coll., *Phys. Rev.* **D55**, 1280 (1997).





Fig. 1. The diagrams contributing to the gluon-photon production of $(\bar{b}c)$ -quarkonium in the leading order of perturbative theory. The c -, b -quarks, gluons and photons are denoted by the thin, thick, spiral and wavy lines, respectively.

$\sigma_{g\gamma}^{B_c^{(*)}}$, pb



$\sqrt{s_{g\gamma}}$, GeV

Fig. 2. The dependence of the gluon-photon cross-section for the production of B_c^* (solid triangle) and B_c (empty triangle) in comparison with the predictions of the fragmentation mechanism (solid and dashed curves, respectively).

$d\sigma_{g\gamma}^{B_c}/dz$, pb




Fig. 3. The $d\sigma_{g\gamma}^{B_c}/dz$ for the B_c meson moving in the gluon hemisphere (dashed histogram) and for the one moving in the photon hemisphere (dotted histogram) at the energy of gluon-photon interaction equal to 100 GeV. For the sake of comparison, the prediction of the fragmentation model is also presented (solid curve).

$$d\sigma_{g\gamma}^{B_c}/dp_T, \text{ pb/GeV}$$




Fig. 4. The differential cross-section for the gluon-photon production of B_c versus the transverse momentum at the energy of 100 GeV. The notations are the same as in Fig. 3.

$$\sigma^{-1} d\sigma_{ep}^{B_c^*}/dy$$





Fig. 5. The differential cross-section normalized to unit area over the rapidity of B_c^* produced in ep -interactions at $\sqrt{s} = 314$ GeV (solid histogram). For the sake of comparison, the normalized distribution with the constant matrix element is shown by the dashed histogram. The rapidity y is defined in the c.m.s. of the colliding beams.

$$\sigma^{-1} d\sigma_{ep}^{B_c^*}/dy(p_T > 30 \text{ GeV})$$



y

Fig. 6. The same as in Fig. 5, but with $p_T > 30 \text{ GeV}$.

$$\sigma^{-1}d\sigma_{ep}^{B_c^*}/dy$$




Fig. 7. The same as in Fig. 5, but with the energy of ep -interactions, $E = 660$ GeV.

$$\sigma^{-1}d\sigma_{ep}^{D_s^*}/dy$$




Fig. 8. The differential cross-section normalized to unit over the rapidity of D_s^* produced in ep -interactions at $\sqrt{s} = 320$ GeV (solid histogram). For the sake of comparison, the normalized distribution with the constant matrix element is shown by the dashed histogram.